

- **Name:** Koichi Goto
 - **Current Position & Affiliation:** Chief, Department of Thoracic Oncology,
National Cancer Center Hospital East
 - **Country:** Japan
-

- **Educational Background:**

March 1990 MD, Kumamoto University School of Medicine
January 2006 PhD, Kumamoto University School of Medicine

- **Professional Experience:**

Recent 2 years experiences are;
Chief, Department of Thoracic Oncology, National Cancer Center Hospital East
Chief, Supportive Care Center, National Cancer Center Hospital East

- **Professional Organizations:**

American Society of Clinical Oncology
Japanese Society of Medical Oncology
The Japan Lung Cancer Society
The Japanese Cancer Association

- **Main Scientific Publications:**

1. Drilon A, Siena S, Dziadziuszko R, Barlesi F, Krebs MG, Shaw AT, de Braud F, Rolfo C, Ahn MJ, Wolf J, Seto T, Cho BC, Patel MR, Chiu CH, John T, Goto K, Karapetis CS, Arkenau HT, Kim SW, Ohe Y, Li YC, Chae YK, Chung CH, Otterson GA, Murakami H, Lin CC, Tan DSW, Prenen H, Riehl T, Chow-Maneval E, Simmons B, Cui N, Johnson A, Eng S, Wilson TR, Doebele RC. Entrectinib in ROS1 fusion-positive non-small-cell lung cancer: integrated analysis of three phase 1-2 trials. *Lancet Oncol*, 2020, 21(2):261-70
2. Naito T, Udagawa H, Sato J, Horinouchi H, Murakami S, Goto Y, Kanda S, Fujiwara Y, Yamamoto N, Zenke Y, Kirita K, Matsumoto S, Yoh K, Niho S, Motoi N, Ohe Y, Ishii G, Goto K. A Minimum Of 100 Tumor Cells in a Single Biopsy Sample Is Required to Assess Programmed Cell Death Ligand 1 Expression in Predicting Patient Response to Nivolumab Treatment in Nonsquamous Non-Small Cell Lung Carcinoma. *J Thorac Oncol*, 2019, 14(10):1818-27
3. Wu YL, Yang JC, Kim DW, Lu S, Zhou J, Seto T, Yang JJ, Yamamoto N, Ahn MJ, Takahashi T, Yamanaka T, Kemner A, Roychowdhury D, Paolini J, Usari T, Wilner KD, Goto K. Phase II Study of Crizotinib in East Asian Patients With ROS1-Positive Advanced Non-Small-Cell Lung Cancer. *J Clin Oncol*, 2018, 36(14):1405-11
4. Nakaoku T, Kohno T, Araki M, Niho S, Chauhan R, Knowles PP, Tsuchihara K,

- Matsumoto S, Shimada Y, Mimaki S, Ishii G, Ichikawa H, Nagatoishi S, Tsumoto K, Okuno Y, Yoh K, McDonald NQ, Goto K. A secondary RET mutation in the activation loop conferring resistance to vandetanib. *Nat Commun*, 2018, 9(1):625
5. Takahashi T, Yamanaka T, Seto T, Harada H, Nokihara H, Saka H, Nishio M, Kaneda H, Takayama K, Ishimoto O, Takeda K, Yoshioka H, Tachihara M, Sakai H, Goto K, Yamamoto N. Prophylactic cranial irradiation versus observation in patients with extensive-disease small-cell lung cancer: a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol*, 2017, 18(5):663-671
6. Hida T, Nokihara H, Kondo M, Kim Y, H, Azuma K, Seto T, Takiguchi Y, Nishio M, Yoshioka H, Imamura F, Hotta K, Watanabe S, Goto K, Satouchi M, Kozuki T, Shukuya T, Nakagawa K, Mitsudomi T, Yamamoto N, Asakawa T, Asabe R, Tanaka T, Tamura T. Alectinib versus crizotinib in patients with ALK-positive non-small-cell lung cancer (J-ALEX): an open-label, randomised phase 3 trial. *Lancet*, 2017, 390(10089):29-39
7. Yoh K, Seto T, Satouchi M, Nishio M, Yamamoto N, Murakami H, Nogami N, Matsumoto S, Kohno T, Tsuta K, Tsuchihara K, Ishii G, Nomura S, Sato A, Ohtsu A, Ohe Y, Goto K. Vandetanib in patients with previously treated RET-rearranged advanced non-small-cell lung cancer (LURET): an open-label, multicentre phase 2 trial. *Lancet Respir Med*, 2017, 5(1):42-50
8. Goto K, Endo M, Kusumoto M, Yamamoto N, Ohe Y, Shimizu A, Fukuoka M. Bevacizumab for non-small-cell lung cancer: A nested case control study of risk factors for hemoptysis. *Cancer Sci*, 2016, 107(12):1837-1842
9. Suzuki K, Yamanaka T, Hashimoto H, Shimada Y, Arata K, Matsui R, Goto K, Takiguchi T, Ohyanagi F, Kogure Y, Nogami N, Nakao M, Takeda K, Azuma K, Nagase S, Hayashi T, Fujiwara K, Shimada T, Seki N, Yamamoto N. Randomized, double-blind, phase III trial of palonosetron versus granisetron in the triplet regimen for preventing chemotherapy-induced nausea and vomiting after highly emetogenic chemotherapy: TRIPLE study. *Ann Oncol*, 2016, 27(8):1601-6
10. Goto K, Ohe Y, Shibata T, Seto T, Takahashi T, Nakagawa K, Tanaka H, Takeda K, Nishio M, Mori K, Satouchi M, Hida T, Yoshimura N, Kozuki T, Imamura F, Kiura K, Okamoto H, Sawa T, Tamura T, for the JCOG0605 investigators. Combined chemotherapy with cisplatin, etoposide, and irinotecan versus topotecan alone as second-line treatment for patients with sensitive relapsed small-cell lung cancer (JCOG0605): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol*, 2016, 17:1147-57